

9FM0/3B: Further Statistics 01 Mark scheme

| Question | Scheme | Marks | AOs |
|------------------|--|-------|------|
| 1 | H_0 : Drivers are equally likely to be recorded speeding on any day of the week H_1 : Drivers are not equally likely to be recorded speeding on any day of the week | B1 | 2.1 |
| | Expected frequency = $\left[\frac{35 + 30 + 28 + 24 + 40 + 51 + 37}{7} \right]$ | M1 | 3.4 |
| | = 35 | A1 | 1.1b |
| | Test statistic = $\frac{(35-35)^2}{35} + \frac{(30-35)^2}{35} + \frac{(28-35)^2}{35} + \dots$ | M1 | 1.1b |
| | = 13.714... | A1 | 1.1b |
| | $\nu = 7 - 1 = 6$ | B1 | 1.1b |
| | $\chi^2_{(6,0.05)} = 12.592$ | B1 | 1.1a |
| | In critical region, sufficient evidence to reject H_0 , Significant evidence at 5% level of significance to reject Jeremy's belief. | A1 | 3.5a |
| (8 marks) | | | |
| Notes | | | |
| | 1 st B1 Both hypotheses correct (condone reference to discrete uniform distribution) 1 st M1 Using uniform model to calculate expected frequencies 1 st A1 35 2 nd M1 Attempting to find $\sum \frac{(O_i - E_i)^2}{E_i}$ or $\sum \frac{O_i^2}{E_i} - N$ may be implied by awrt 13.7 2 nd A1 awrt 13.7 2 nd B1 Degrees of freedom = 6 may be implied by a correct CV 3 rd B1 awrt 12.6 3 rd A1 Evaluating the outcome of a model by drawing correct inference in context | | |

| Question | Scheme | Marks | AOs |
|------------------|---|------------|------|
| 2(a) | $\text{Var}(Y) = E(Y^2) - [E(Y)]^2$ | M1 | 3.1a |
| | $E(Y) = \frac{1}{2}a + 2 \times \frac{3}{10} + 7 \times \frac{1}{5} [= \frac{1}{2}a + 2]$ | B1 | 1.1b |
| | $E(Y^2) = \frac{1}{2}a^2 + 4 \times \frac{3}{10} + 49 \times \frac{1}{5} [= \frac{1}{2}a^2 + 11]$ | B1 | 2.1 |
| | $28 = \frac{1}{2}a^2 + 11 - (\frac{1}{2}a + 2)^2$ | M1 | 1.1b |
| | $\frac{1}{4}a^2 - 2a - 21 = 0 \rightarrow a = \dots$ | M1 | 1.1b |
| | $a = -6$ since $E(Y) < 0$ | A1 | 2.2a |
| | | (6) | |
| (b) | $\left(\frac{1}{3 - (-6)}\right) \times \frac{1}{2} + \left(\frac{1}{3 - 2}\right) \times \frac{3}{10} + \left(\frac{1}{3 - 7}\right) \times \frac{1}{5}$ | M1 | 2.1 |
| | $= \frac{11}{36}$ | A1ft | 1.1b |
| | | (2) | |
| (8 marks) | | | |
| Notes | | | |
| (a) | 1 st M1 Realising that $\text{Var}(Y) = E(Y^2) - [E(Y)]^2$ is required 1 st B1 Correct expression for $E(Y)$ 2 nd B1 Correct expression for $E(Y^2)$ 2 nd M1 Equating their expression for $\text{Var}(Y) = 28$ 3 rd M1 Solving the equation to find at least 1 value of a A1 -6 only | | |
| (b) | M1 Correct expression for $E\left(\frac{1}{3 - Y}\right)$ or for finding all values of $\frac{1}{3 - Y}$ A1ft $\frac{11}{36}$ or awrt 0.306 ft on $a < -4$ | | |

| Question | Scheme | Marks | AOs |
|-------------------|---|------------|------|
| 3(a) | $W \sim \text{Po}(0.45n)$ | M1 | 3.1b |
| | $[P(W = 0) =] e^{-0.45n} < 0.05$ | M1 | 1.1b |
| | $n > 6.657\dots$ | | |
| | $n = 7$ | A1 | 1.1b |
| | | (3) | |
| (b) | $X \sim \text{Po}(5 \times 0.45 + 5 \times 0.2)$ [$\text{Po}(3.25)$] | M1 | 3.3 |
| | $P(X = 2) = 0.20478\dots$ awrt <u>0.205</u> | A1 | 1.1b |
| | The model is only valid if Tim and Sue make errors independently | B1 | 3.5b |
| | | (3) | |
| (c) | $P(X = 0) = 0.03877\dots$ | M1 | 3.1b |
| | $Y \sim B(10, '0.03877\dots')$ | M1 | 3.3 |
| | $P(Y \geq 2) = 1 - P(Y \leq 1)$ | M1 | 1.1b |
| | = awrt <u>0.055</u> | A1 | 1.1b |
| | | (4) | |
| (10 marks) | | | |
| Notes | | | |
| (a) | 1 st M1 Understanding that a $P(0.45n)$ model is required here 2 nd M1 For correct inequality A1 $n = 7$ cao | | |
| (b) | M1 Setting up a combined Po model A1 awrt 0.205 B1 Understanding that model is only valid if the two parts are independent | | |
| (c) | 1 st M1 For using Poisson distribution 2 nd M1 Setting up binomial distribution 3 rd M1 For finding $1 - P(Y \leq 1)$ from binomial 2 nd A1 awrt 0.055 | | |

| Question | Scheme | Marks | AOs |
|-------------------|---|----------|--------------|
| 4(a) | $n = 2$ and $p = 0.6$ | B1 B1 | 1.1b 1.1b |
| | | (2) | |
| (b)(i) | $P(X=1) = \text{coefficient of } t \quad G_X(t) = 0.16 + 0.48t + 0.36t^2$ | M1 | 1.1b |
| | $P(X=1) = \underline{0.48}$ | A1 | 1.1b |
| | | (2) | |
| (ii) | $E(X) = G'_X(1)$ | M1 | 2.1 |
| | $G'_X(t) = 2(0.4 + 0.6t) \times 0.6$ | M1 | 1.1b |
| | $G'_X(1) = 1.2$ | A1 | 1.1b |
| | | (3) | |
| (c) | $G_Y(t) = G_X(t) \times G_X(t)$ | | |
| | $G_Y(t) = (0.4 + 0.6t)^4$ | B1 | 3.1a |
| | $G'_Y(t) = 4(0.4 + 0.6t)^3 \times 0.6$ | M1 | 2.1 |
| | $G'_Y(1) = 2.4$ | A1 | 1.1b |
| | $G''_Y(t) = 7.2(0.4 + 0.6t)^2 \times 0.6$ | M1 | 2.1 |
| | $G''_Y(1) = 4.32$ | A1 | 1.1b |
| | $E(Y^2) [= \text{Var}(Y) + [E(Y)]^2] = G'_Y(1) + G''_Y(1)$ | M1 | 1.1b |
| | $E(Y^2) = 2.4 + 4.32 = 6.72^*$ | A1*cso | 1.1b |
| | | (7) | |
| (14 marks) | | | |
| Notes | | | |
| (a) | 1 st B1 $n = 2$ 2 nd B1 $p = 0.6$ | | |
| (b)(i) | M1 Finding coefficient of t A1 0.48oe | | |
| (b)(ii) | 1 st M1 Realising $G'_X(1)$ is needed 2 nd M1 Differentiation A1 1.2cao | | |
| (c) | B1 Correct use of $G_Y(t) = G_X(t) \times G_X(t)$ 1 st M1 Differentiation to find $G'_Y(t)$ 1 st A1 $G'_Y(1) = 2.4$ 2 nd M1 Differentiation to find $G''_Y(t)$ 2 nd A1 $G''_Y(1) = 4.32$ 3 rd M1 Realising $E(Y^2) = G'_Y(1) + G''_Y(1)$ 3 rd A1*cso 6.72 | | |

| Question | Scheme | Marks | AOs |
|------------------|---|-------|------|
| 5(a) | $H_0 : p = 0.03$ $H_1 : p < 0.03$ | B1 | 2.5 |
| | $X \sim \text{Geo}(0.03)$ | M1 | 3.3 |
| | $P(X \geq c) < 0.05$ $(1 - 0.03)^{c-1} < 0.05$ | M1 | 3.4 |
| | $c - 1 > \frac{\log 0.05}{\log 0.97}$ | M1 | 1.1b |
| | $c > 99.35\dots$ critical region $X \geq 100$ | A1 | 2.2a |
| | | (5) | |
| (b) | $P(X \geq 100) = 0.97^{99}$ | M1 | 3.4 |
| | = <u>0.0490</u> | A1 | 1.1b |
| | | (2) | |
| (c) | Critical region $X \geq 100$ 94 is not in the critical region [$P(X \geq 94) = 0.0588\dots > 0.05$] | M1 | 1.1b |
| | Do not reject H_0 There is insufficient evidence at the 5% level of significance that the proportion of visitors making a purchase is less than 0.03 | A1 | 2.2b |
| | | (2) | |
| (9 marks) | | | |
| Notes | | | |
| (a) | B1 Both hypotheses correct using correct notation 1 st M1 Realising that the model Geo (0.03) is needed. May be implied by its use 2 nd M1 Using the model to find an expression for $P(X \geq c)$ 3 rd M1 Finding a valid method to solve the inequality A1 Correct critical region | | |
| (b) | M1 Using Geo(0.03) model with 100 A1 0.049 or awrt 0.0490 | | |
| (c) | M1 Comparing 94 with their critical value A1 Fully correct solution and drawing a correct inference in context. | | |

| Question | Scheme | Marks | AOs |
|-------------------|---|------------|-------------|
| 6(a) | P(Type I error) = <u>0.05</u> | B1 | 1.2 |
| | | (1) | |
| (b) | $\bar{X} \sim N(120, \frac{3^2}{10}) \quad P(\bar{X} > c) < 0.05$ | M1 | 3.1b |
| | $\frac{c-120}{\frac{3}{\sqrt{10}}} > 1.6449$ | M1 | 3.4 |
| | $c > 121.56\dots$ | A1 | 1.1b |
| | $P(\bar{X} > 121.56 \mid \mu = 122)$ | | |
| | $P\left(Z > \frac{121.56 - 122}{\frac{3}{\sqrt{10}}}\right) = P(Z > -0.4638\dots)$ | M1 | 2.1 |
| | $= 0.6786\dots \quad = 0.68*(2sf)$ | A1*cso | 1.1b |
| | | (5) | |
| (c) | Power of Alex's test is smaller than power of Gizel's test since the null hypothesis is less likely to be rejected/Type II error has increased. | B1 B1 | 2.2a 2.4 |
| | | (2) | |
| (d) | $\frac{c-120}{\frac{3}{\sqrt{n}}} > 1.6449$ | M1 | 3.4 |
| | $c > 120 + 1.6449 \times \frac{3}{\sqrt{n}}$ | A1 | 1.1b |
| | $P(\bar{X} > c \mid \mu = 122) > 0.9$ | | |
| | $\frac{(120 + 1.6449 \times \frac{3}{\sqrt{n}}) - 122}{\frac{3}{\sqrt{n}}} < -1.2816$ | M1 | 2.1 |
| | $2.9265 \frac{3}{\sqrt{n}} < 2 \quad \rightarrow \quad \sqrt{n} > 4.38\dots$ | M1 | 1.1b |
| | $n > 19.26\dots \quad n = \underline{\underline{20}}$ | A1 | 1.1b |
| | | (5) | |
| (e) | (As they both have the same size/Type I error and) Joseph's test has a higher power, so Joseph's test is recommended. | M1 A1 | 2.4 2.2b |
| | | (2) | |
| (15 marks) | | | |
| Notes | | | |
| (a) | B1 0.05oe | | |
| (b) | 1 st M1 Selecting correct normal model | | |
| | 2 nd M1 Using model to standardise and set up inequality | | |
| (c) | 1 st A1 Correct critical region | | |
| | 3 rd M1 Correct probability statement to find power | | |
| (d) | 2 nd A1*cso awrt 0.68 with no errors seen. | | |
| | B1 Correct deduction about the size of the two tests | | |
| (e) | B1 Correct explanation | | |

| | |
|------------|--|
| (d) | 1 st M1 Using normal model to find critical region |
| | 1 st A1 Correct critical region in terms of n |
| | 2 nd M1 Setting up comparison with $ 1.2816 $ to find n |
| | 3 rd M1 Solving equation to $\sqrt{n} > \dots$ |
| (e) | 2 nd A1 20cao |
| | M1 Comparison of powers A1 Correct conclusion based on power |

| Question | Scheme | Marks | AOs |
|-------------------|--|------------|--------------|
| 7(a) | $[X \sim \text{NB}(12, \frac{3}{4})]$ | | |
| | $\binom{14}{11} \times \left(\frac{3}{4}\right)^{12} \times \left(\frac{1}{4}\right)^3$ | M1 | 3.3 |
| | = awrt 0.180 | A1 | 1.1b |
| | | (2) | |
| (b) | $P(X > 13) = 1 - [P(X = 12) + P(X = 13)]$ | B1 | 3.1b |
| | $1 - \left(\left(\frac{3}{4}\right)^{12} + \binom{12}{11} \left(\frac{3}{4}\right)^{12} \times \left(\frac{1}{4}\right) \right)$ | M1 | 1.1b |
| | = awrt 0.873 | A1 | 1.1b |
| | | (3) | |
| (c) | $E(X) = \frac{12}{\frac{3}{4}} = 16$ | M1 | 3.1b |
| | $\text{Var}(X) = \frac{12(\frac{1}{4})}{(\frac{3}{4})^2} = \frac{16}{3}$ | A1 | 1.1b |
| | $\bar{X} \square N\left(16, \frac{16}{30} (= 0.1\dot{7})\right)$ | M1 A1ft | 3.1b 1.1b |
| | $P(\bar{X} > 15.5) = P\left(Z > \frac{15.5 - 16}{\sqrt{0.1\dot{7}}}\right)$ | M1 | 3.4 |
| | = $P(Z > -1.1858\dots)$ | | |
| | = awrt 0.882/0.883 | A1 | 1.1b |
| | | | (6) |
| (11 marks) | | | |
| Notes | | | |
| (a) | M1 Selecting correct model: negative binomial or $B(14, \frac{3}{4})$ with extra success A1 0.18 or awrt 0.180 | | |
| (b) | B1 Realising that $P(X > 13) = 1 - [P(X = 12) + P(X = 13)]$ M1 Correct form using negative binomial A1 awrt 0.873 | | |
| (c) | 1 st M1 Realising that both the mean and variance of NB are required 1 st A1 Both mean and variance correct (may be implied by correct standardisation) 2 nd M1 Using CLT to model $\bar{X} \sim N('16', \dots)$ 2 nd A1ft Fully correct (or correct ft) normal distribution model for \bar{X} 3 rd M1 Using the normal model to find $P(\bar{X} > 15.5)$. Can be awarded for correct (ft) standardisation 3 rd A1 awrt 0.882 or 0.883 | | |